Search Topics

Webcasts

Science Links

Contact Us

Our Scientists

UCSB Science Line



Ernest Rutherford bombarded nitrogen atoms with helium nuclei (alpha particles), and hydrogen atoms (protons) were formed as a result. From this, he concluded that nitrogen nuclei contained protons.

Note from ScienceLine moderator:

Please watch this <u>video</u> as the best way to illustrate the Rutherford's experiment for the discovery of the proton.

Answer 2:

Great question! Rutherford's experiment was incredibly powerful due to its simplicity, use of cheap materials, and the information he uncovered. Rutherford's main focus was on studying the nucleus through radioactive decay.

After the electron was discovered by Thomson in 1897 and after Rutherford's work on discovering the existence of the nucleus 1911, it was known that there must exist particles of positive charge to balance the negatively charged electrons to create electrically neutral atoms.

The series of experiments, performed by Rutherford and his student James Chadwick, **consisted of changing one element into another by hitting atoms with high energy alpha particles.** Specifically, they noticed that nitrogen, oxygen, and aluminum, when hit with an alpha particle, disintegrated and emitted a fast particle of positive charge. Or said more specifically, hydrogen nuclei were always emitted in the process. In a dark room, they were able to observe flashes of light when alpha particles hit the target. Alpha particles, which were also discovered by Rutherford, are spontaneously emitted by radioactive materials such as uranium. It was realized that the positive charge of any nucleus could be accounted for by a whole (integer) number of positively charged hydrogen nuclei, which were named protons by Rutherford in 1920.

They immediately wondered what was left behind on the target after this process occurred. The conclusion was that the target captured the alpha particle (2+ charge) and emitted a proton (1+ charge), resulting in the target having a nuclear charge different than before. The target was now an isotope of another element. In the case of a nitrogen target, the nitrogen had a nuclear charge of 8 instead of seven, making it an isotope of oxygen (an isotope is an atom with the same charge but a different atomic weight.) This describes how the proton itself was discovered.

Many key experiments by Rutherford, his students, and others occurred before this and led to the understanding of the atom that is still the accepted model today.

Click <u>Here</u> to return to the search form.





National Science Foundation

This program is co-sponsored by the National Science Foundation and UCSB School-University Partnerships Copyright © 2017 The Regents of the University of California, All Rights Reserved. <u>UCSB Terms of Use</u>